

2014-2015 South Carolina Adult Tobacco Survey

Division of Tobacco Prevention and Control
South Carolina Department of Health and Environment Control



Contents

| | |
|--|----|
| Table 1: Percentage of SC adults who reported ever ¹ smoking cigarettes and currently ² smoking cigarettes by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. | 3 |
| Table 2: Percentage of South Carolina adult current ¹ smokers by smoking frequency (current everyday & someday smokers) – Results from the 2014-2015 SC Adult Tobacco Survey. | 4 |
| Table 3: Percentage of South Carolina current ¹ adult smokers by average number of cigarettes smoked per day – Results from the 2012 SC Adult Tobacco Survey..... | 4 |
| Table 4: Percentage of SC adults who reported ever ¹ smoking e-cigarettes and currently ² smoking e-cigarettes by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. | 5 |
| Table 5: Percentage of SC adults who reported current dual use ¹ of e-cigarettes ² and conventional cigarettes ³ or any tobacco product ⁴ by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. | 7 |
| Table 6: Percentage of South Carolina adults who reported ever ¹ using other tobacco products (cigars, smokeless tobacco, water pipe, snus) by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. | 9 |
| Table 7: Percentage of South Carolina adults who report current use of menthol cigarettes ¹ and other tobacco products ² (cigars, smokeless tobacco, water pipe, regular pipe, and snus), by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. | 11 |
| Table 8: Percentage of South Carolina adults exposed ¹ to Secondhand Smoke (SHS) at various locations – Results from the 2014-2015 SC Adult Tobacco Survey. | 13 |
| Table 9: Percentage of South Carolina adult current smokers ¹ and adults exposed to Secondhand Smoke (SHS) ² at various locations among those with chronic conditions ³ – Results from the 2014-2015 SC Adult Tobacco Survey..... | 13 |
| Table 10: Percentage of South Carolina adults who reported being screened ¹ for tobacco use by a health care provider during their visit in the past 12 months, overall and by chronic condition ² status – Results from the 2014-2015 SC Adult Tobacco Survey..... | 14 |
| Table 11: Percentage of South Carolina adults who reported receiving smoking cessation help from their health care provider during their visit in the past 12 months – Results from the 2014-2015 SC Adult Tobacco Survey..... | 15 |
| Table 12: Percentage of South Carolina adults who reported their quit intentions, awareness of cessation services and use of cessation services during their quit attempt – Results from the 2014-2015 SC Adult Tobacco Survey..... | 15 |
| Table 13: Percentage of South Carolina adult current smokers who reported having quit for one or more days in the past 12 months, and sustained their abstinence – Results from the 2014-2015 SC Adult Tobacco Survey..... | 15 |
| Table 14: Percentage of South Carolina adults who reported having seen one or more ads about not smoking, by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. | 16 |

Table 15: Percentage of South Carolina adults who stated that the ads seen contributed to their attempt to quit, by demographics – Results from the 2014-2015 SC Adult Tobacco Survey. 17

Table 16: Percentage of South Carolina adults who mentioned that smoking is never allowed at various locations, overall and by current smoking status – Results from the 2014-2015 SC Adult Tobacco Survey. 18

Table 17: Percentage of South Carolina adults who thought smoking should never be allowed at various locations, overall and by current smoking status – Results from the 2014-2015 SC Adult Tobacco Survey. 18

Table 18: Percentage of South Carolina adults who support a cigarette tax increase – Results from the 2014-2015 SC Adult Tobacco Survey. 18

Table 1: Percentage of SC adults who reported ever¹ smoking cigarettes and currently² smoking cigarettes by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | Total Respondents | Ever Use of Cigarettes | | Current Use of Cigarettes | |
|--|-------------------|------------------------|---------------------|---------------------------|---------------|
| | | % | 95% CI ³ | % | 95% CI |
| TOTAL | 7503 | 43.1 | (±1.7) | 19.0 | (±1.5) |
| Gender | | | | | |
| Male | 3072 | 49.8 | (±2.6) | 21.3 | (±2.4) |
| Female | 4431 | 36.8 | (±2.2) | 16.9 | (±2.0) |
| Age Group | | | | | |
| 18-24 | 429 | 22.8 | (±5.1) | 15.6 | (±4.3) |
| 25-34 | 659 | 44.9 | (±5.5) | 29.5 | (±5.4) |
| 35-44 | 752 | 39.4 | (±4.9) | 21.0 | (±4.4) |
| 45-54 | 1146 | 45.6 | (±3.8) | 24.3 | (±3.5) |
| 55-64 | 1642 | 50.5 | (±3.2) | 18.1 | (±2.6) |
| 65+ | 2697 | 50.2 | (±2.4) | 8.4 | (±1.4) |
| Race/Ethnicity | | | | | |
| non-Hispanic White | 4827 | 47.1 | (±2.1) | 18.8 | (±1.9) |
| non-Hispanic Black | 1817 | 34.1 | (±3.2) | 18.7 | (±2.9) |
| non-Hispanic Asian | 35 | 13.7 | (±13.3) | 4.8 | (±4.8) |
| non-Hispanic American Indian & Alaska Native | 38 | 63.5 | (±17.5) | 48.9 | (±19.4) |
| non-Hispanic Other | 436 | 60.7 | (±7.1) | 37.8 | (±7.8) |
| Hispanic | 131 | 28.8 | (±12.5) | 17.4 | (±11.2) |
| Education | | | | | |
| < High School | 706 | 57.4 | (±5.8) | 35.2 | (±5.9) |
| High School or GED | 2014 | 46.0 | (±3.2) | 22.1 | (±2.7) |
| Some College | 2255 | 41.6 | (±2.9) | 17.3 | (±2.5) |
| College Degree or Higher | 2475 | 31.2 | (±2.4) | 6.0 | (±1.3) |
| Marital Status | | | | | |
| Married | 3854 | 45.3 | (±2.3) | 14.7 | (±2.0) |
| Not Married | 3568 | 40.4 | (±2.6) | 23.4 | (±2.4) |
| Household Poverty Status | | | | | |
| Below 100% FPL | 1209 | 44.5 | (±5.0) | 23.0 | (±5.1) |
| At or Above 100% FPL | 3304 | 43.6 | (±2.7) | 14.5 | (±2.3) |
| Unspecified | 2990 | 41.9 | (±2.4) | 22.3 | (±2.1) |
| Residence | | | | | |
| Urban | 5489 | 43.6 | (±2.0) | 19.4 | (±1.8) |
| Rural | 1854 | 41.3 | (±3.6) | 17.5 | (±2.8) |
| DHEC Region | | | | | |
| Upstate | 1782 | 43.0 | (±3.4) | 19.4 | (±3.1) |
| Pee Dee | 1782 | 44.1 | (±3.3) | 20.2 | (±2.9) |
| Midlands | 2062 | 44.6 | (±3.3) | 19.3 | (±3.0) |
| Low Country | 1717 | 39.4 | (±3.2) | 16.2 | (±2.6) |

- 1 - To determine ever use, respondents were asked ‘Have you ever tried cigarette smoking, even one or two puffs?’
- 2 - To determine current use, respondents were asked two questions ‘Have you smoked at least 100 cigarettes in your entire life?’ and ‘Do you now smoke cigarettes every day, some days, or not at all? Those who smoked at least 100 cigarettes in lifetime and currently smoke every day or someday are coded as current smokers.
- 3 - Ninety-five percent confidence interval. Note large confidence intervals for distribution by race/ethnicity

Table 2: Percentage of South Carolina adult current¹ smokers by smoking frequency (current everyday & someday smokers) – Results from the 2014-2015 SC Adult Tobacco Survey.

| Smoking Frequency | Current Smokers (N=1097) | |
|-------------------|--------------------------|---------------------|
| | % | 95% CI ² |
| Everyday Smoker | 74.2 | (±3.7) |
| Someday Smoker | 25.8 | (±3.7) |

- 1 - To determine current use, respondents were asked two questions ‘Have you smoked at least 100 cigarettes in your entire life?’ and ‘Do you now smoke cigarettes every day, some days, or not at all? Those who smoked at least 100 cigarettes in lifetime and currently smoke every day or someday are coded as current smokers.
- 2 - Ninety-five percent confidence interval.

Table 3: Percentage of South Carolina current¹ adult smokers by average number of cigarettes smoked per day – Results from the 2012 SC Adult Tobacco Survey.

| Average Number of Cigarettes Smoked Per Day (CPD) | All Current Smokers (N=1097) | | Current Everyday Smokers (N=788) | | Current Someday Smokers (N=309) | |
|---|------------------------------|---------------------|----------------------------------|--------|---------------------------------|--------|
| | % | 95% CI ² | % | 95% CI | % | 95% CI |
| 1-10 CPD | 53.1 | (±4.9) | 40.9 | (±5.7) | 91.2 | (±4.6) |
| 11-20 CPD | 39.0 | (±4.8) | 49.5 | (±5.8) | 6.5 | (±3.9) |
| >20 CPD | 7.8 | (±2.1) | 9.6 | (±2.6) | 2.2 | (±2.2) |

- 1 - To determine current use, respondents were asked two questions ‘Have you smoked at least 100 cigarettes in your entire life?’ and ‘Do you now smoke cigarettes every day, some days, or not at all? Those who smoked at least 100 cigarettes in lifetime and currently smoke every day or someday are coded as current smokers.
- 2 - Ninety-five percent confidence interval.

Table 4: Percentage of SC adults who reported ever¹ smoking e-cigarettes and currently² smoking e-cigarettes by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | Ever Tried e-Cigarettes | | | Current Use of e-Cigarettes | | |
|--|-------------------------|-------------|---------------------|-----------------------------|-----------------|---------------|
| | Total Respondents | % | 95% CI ³ | Total Respondents | % | 95% CI |
| TOTAL | 7497 | 16.2 | (±1.5) | 7492 | 6.2 | (±1.1) |
| Gender | | | | | | |
| Male | 3070 | 18.2 | (±2.4) | 3068 | 7.3 | (±1.9) |
| Female | 4427 | 14.4 | (±1.8) | 4424 | 5.1 | (±1.2) |
| Age Group | | | | | | |
| 18-24 | 429 | 28.6 | (±5.7) | 428 | 11.5 | (±4.0) |
| 25-34 | 657 | 28.0 | (±5.2) | 657 | 11.8 | (±4.5) |
| 35-44 | 752 | 17.1 | (±4.1) | 751 | 7.3 | (±2.8) |
| 45-54 | 1145 | 14.1 | (±2.6) | 1146 | 4.1 | (±1.6) |
| 55-64 | 1641 | 11.4 | (±2.2) | 1640 | 3.6 | (±1.3) |
| 65+ | 2695 | 4.9 | (±1.2) | 2693 | 1.8 | (±0.8) |
| Race/Ethnicity | | | | | | |
| non-Hispanic White | 4825 | 17.7 | (±1.8) | 4820 | 6.6 | (±1.3) |
| non-Hispanic Black | 1815 | 11.2 | (±2.5) | 1814 | 3.5 | (±1.6) |
| non-Hispanic Asian | 35 | 2.7 | (±2.7) | 35 | -- ⁴ | -- |
| non-Hispanic American Indian & Alaska Native | 38 | 27.8 | (±19.7) | 38 | 12.3 | (±12.3) |
| non-Hispanic Other | 436 | 35.5 | (±7.8) | 435 | 16.9 | (±7.4) |
| Hispanic | 131 | 14.9 | (±11.3) | 131 | 10.7 | (±10.7) |
| Education | | | | | | |
| < High School | 702 | 21.8 | (±5.0) | 701 | 7.9 | (±3.2) |
| High School or GED | 2014 | 19.0 | (±2.9) | 2011 | 8.4 | (±2.3) |
| Some College | 2254 | 17.6 | (±2.6) | 2253 | 6.2 | (±2.0) |
| College Degree or Higher | 2474 | 6.7 | (±1.4) | 2474 | 2.4 | (±0.8) |
| Marital Status | | | | | | |
| Married | 3853 | 12.9 | (±1.9) | 3853 | 4.9 | (±1.5) |
| Not Married | 3563 | 19.7 | (±2.2) | 3559 | 7.6 | (±1.6) |
| Household Poverty Status | | | | | | |
| Below 100% FPL | 1206 | 10.9 | (±3.2) | 1205 | 5.3 | (±2.7) |
| At or Above 100% FPL | 3303 | 13.6 | (±2.5) | 3300 | 5.8 | (±1.9) |
| Unspecified | 2988 | 21.2 | (±2.1) | 2987 | 7.0 | (±1.3) |
| Residence | | | | | | |
| Urban | 5484 | 17.2 | (±1.7) | 5483 | 6.6 | (±1.3) |
| Rural | 1853 | 11.6 | (±2.5) | 1850 | 4.3 | (±1.9) |
| DHEC Region | | | | | | |
| Upstate | 1782 | 16.7 | (±3.1) | 1776 | 7.2 | (±2.6) |
| Pee Dee | 1779 | 16.2 | (±2.8) | 1780 | 4.8 | (±1.4) |
| Midlands | 2061 | 15.0 | (±2.7) | 2061 | 5.9 | (±1.9) |
| Low Country | 1715 | 16.8 | (±2.8) | 1716 | 6.0 | (±1.8) |

- 1 - To determine ever use, respondents were asked 'Have you ever used any of the following products: Electronic cigarettes such as "Smoking Everywhere" or "Njoy"?'.
- 2 - To determine current use, respondents were asked 'During the past 30 days, [...], on how many days did you use Electronic cigarettes, such as "Smoking Everywhere" or "Njoy"?' Those who used on at least 1 day are coded as current users.
- 3 - Ninety-five percent confidence interval.
- 4 - Data are statistically unreliable because sample size was <50. Note large confidence intervals for distribution by race/ethnicity.

Table 5: Percentage of SC adults who reported current dual use¹ of e-cigarettes² and conventional cigarettes³ or any tobacco product⁴ by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | Dual Use With Conventional Cigarettes | | | Dual Use With Any Tobacco Product | | |
|--|---------------------------------------|-----------------|---------------------|-----------------------------------|------------|---------------|
| | Total Respondents | % | 95% CI ⁵ | Total Respondents | % | 95% CI |
| TOTAL | 7455 | 4.1 | (±0.9) | 7438 | 4.6 | (±1.0) |
| Gender | | | | | | |
| Male | 3052 | 5.1 | (±1.7) | 3040 | 6.0 | (±1.8) |
| Female | 4403 | 3.2 | (±0.9) | 4398 | 3.3 | (±0.9) |
| Age Group | | | | | | |
| 18-24 | 428 | 6.4 | (±2.9) | 427 | 8.8 | (±3.6) |
| 25-34 | 653 | 7.8 | (±4.0) | 653 | 8.4 | (±4.0) |
| 35-44 | 748 | 4.4 | (±2.5) | 747 | 5.1 | (±2.5) |
| 45-54 | 1146 | 3.5 | (±1.5) | 1143 | 3.5 | (±1.5) |
| 55-64 | 1634 | 2.7 | (±1.1) | 1634 | 2.8 | (±1.1) |
| 65+ | 2676 | 1.3 | (±0.7) | 2663 | 1.3 | (±0.7) |
| Race/Ethnicity | | | | | | |
| non-Hispanic White | 4799 | 3.9 | (±0.9) | 4784 | 4.5 | (±1.0) |
| non-Hispanic Black | 1803 | 2.7 | (±1.5) | 1801 | 2.8 | (±1.6) |
| non-Hispanic Asian | 35 | -- ⁶ | -- | 35 | -- | -- |
| non-Hispanic American Indian & Alaska Native | 38 | 12.3 | (±12.3) | 38 | 12.3 | (±12.3) |
| non-Hispanic Other | 433 | 14.8 | (±7.4) | 433 | 15.8 | (±7.4) |
| Hispanic | 131 | 10.7 | (±10.7) | 131 | 10.7 | (±10.7) |
| Education | | | | | | |
| < High School | 695 | 6.9 | (±3.1) | 690 | 7.6 | (±3.3) |
| High School or GED | 1999 | 4.6 | (±1.6) | 1994 | 5.6 | (±1.8) |
| Some College | 2244 | 4.2 | (±1.9) | 2241 | 4.4 | (±1.9) |
| College Degree or Higher | 2466 | 1.5 | (±0.7) | 2463 | 1.8 | (±0.7) |
| Marital Status | | | | | | |
| Married | 3837 | 3.1 | (±1.3) | 3827 | 3.3 | (±1.3) |
| Not Married | 3540 | 5.2 | (±1.3) | 3532 | 6.0 | (±1.4) |
| Household Poverty Status | | | | | | |
| Below 100% FPL | 1193 | 5.1 | (±2.8) | 1187 | 5.1 | (±2.8) |
| At or Above 100% FPL | 3283 | 3.5 | (±1.6) | 3277 | 3.8 | (±1.7) |
| Unspecified | 2979 | 4.3 | (±1.1) | 2974 | 5.2 | (±1.1) |
| Residence | | | | | | |
| Urban | 5459 | 4.3 | (±1.1) | 5448 | 4.9 | (±1.1) |
| Rural | 1841 | 3.4 | (±1.8) | 1834 | 3.7 | (±1.8) |
| DHEC Region | | | | | | |
| Upstate | 1772 | 5.3 | (±2.4) | 1765 | 6.2 | (±2.6) |
| Pee Dee | 1771 | 3.7 | (±1.2) | 1768 | 3.8 | (±1.2) |
| Midlands | 2051 | 3.6 | (±1.4) | 2047 | 3.7 | (±1.4) |
| Low Country | 1706 | 3.4 | (±1.4) | 1702 | 4.7 | (±1.7) |

1 - Dual use is defined as a respondent's concurrent use of both electronic cigarettes and conventional cigarettes or electronic cigarettes and any tobacco product (cigarettes, cigars, smokeless tobacco, regular pipe, water pipe, snus).

2 - To determine current use of electronic cigarettes, respondents were asked 'During the past 30 days, [...], on how many days did you use Electronic cigarettes, such as "Smoking Everywhere" or "Njoy"?' Those who used on at least 1 day are coded as current users.

3 - To determine current use of conventional cigarettes, respondents were asked two questions 'Have you smoked at least 100 cigarettes in your entire life?' and 'Do you now smoke cigarettes every day, some days, or not at all?' Those who smoked at least 100 cigarettes in lifetime and currently smoke every day or someday are coded as current smokers.

4 - To determine current use of other tobacco products, respondents were asked the following questions: "During the past 30 days, on how many days did you use chewing tobacco, snuff, or dip, such as Skoal, Copenhagen, Grizzly, Levi Garrett, Red Man, or Day's Work?", "During the past 30 days, on how many days did you smoke cigars, cigarillos, or very small cigars that look like cigarettes?", "During the past 30 days, on how many days did you smoke tobacco in a water pipe?", "During the past 30 days, on how many days did you smoke tobacco in a pipe other than a water pipe?", "During the past 30 days, on how many days did you use snus?". Those who used at least one of those tobacco products on 1 or more days are coded as current any tobacco users.

5 - Ninety-five percent confidence interval.

6 - Data are statistically unreliable because sample size was <50. Note large confidence intervals for distribution by race/ethnicity.

Table 6: Percentage of South Carolina adults who reported ever¹ using other tobacco products (cigars, smokeless tobacco, water pipe, snus) by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | Cigars (N=7494) | | Smokeless Tobacco (N=7499) | | Water Pipe (N=7486) | | Snus (N=7498) | |
|--|-----------------|---------------------|----------------------------|---------------|---------------------|---------------|---------------|---------------|
| | % | 95% CI ² | % | 95% CI | % | 95% CI | % | 95% CI |
| TOTAL | 38.2 | (±1.7) | 23.9 | (±1.5) | 9.8 | (±1.2) | 5.0 | (±0.9) |
| Gender | | | | | | | | |
| Male | 57.5 | (±2.7) | 42.2 | (±2.6) | 13.1 | (±2.0) | 9.0 | (±1.7) |
| Female | 20.5 | (±1.8) | 7.1 | (±1.1) | 6.8 | (±1.2) | 1.4 | (±0.5) |
| Age Group | | | | | | | | |
| 18-24 | 33.3 | (±5.8) | 22.4 | (±5.3) | 26.2 | (±5.3) | 7.3 | (±2.9) |
| 25-34 | 42.6 | (±5.3) | 24.3 | (±4.8) | 21.0 | (±4.3) | 8.6 | (±3.6) |
| 35-44 | 38.9 | (±4.8) | 29.1 | (±4.6) | 8.3 | (±2.9) | 7.1 | (±2.5) |
| 45-54 | 40.7 | (±3.8) | 30.3 | (±3.6) | 4.8 | (±1.7) | 5.3 | (±1.7) |
| 55-64 | 40.3 | (±3.1) | 23.3 | (±2.7) | 3.2 | (±1.0) | 2.5 | (±0.9) |
| 65+ | 35.1 | (±2.4) | 16.9 | (±2.0) | 1.3 | (±0.6) | 1.4 | (±0.6) |
| Race/Ethnicity | | | | | | | | |
| non-Hispanic White | 42.5 | (±2.1) | 28.5 | (±2.0) | 9.4 | (±1.3) | 5.9 | (±1.0) |
| non-Hispanic Black | 28.1 | (±3.1) | 13.7 | (±2.3) | 8.2 | (±2.2) | 1.8 | (±1.0) |
| non-Hispanic Asian | 16.4 | (±14.3) | 3.4 | (±3.4) | -- ³ | -- | -- | -- |
| non-Hispanic American Indian & Alaska Native | 41.6 | (±19.6) | 24.4 | (±16.0) | -- | -- | -- | -- |
| non-Hispanic Other | 57.0 | (±7.2) | 32.9 | (±7.4) | 22.9 | (±7.5) | 11.0 | (±4.8) |
| Hispanic | 26.6 | (±11.6) | 15.9 | (±11.3) | 17.1 | (±11.2) | -- | -- |
| Education | | | | | | | | |
| < High School | 35.5 | (±5.5) | 26.5 | (±5.1) | 6.6 | (±3.4) | 3.0 | (±1.3) |
| High School or GED | 39.8 | (±3.2) | 25.2 | (±2.9) | 9.3 | (±2.1) | 5.7 | (±1.6) |
| Some College | 37.2 | (±2.9) | 23.2 | (±2.7) | 11.3 | (±2.3) | 5.9 | (±2.1) |
| College Degree or Higher | 40.1 | (±2.6) | 22.1 | (±2.3) | 10.7 | (±1.9) | 4.4 | (±1.2) |
| Marital Status | | | | | | | | |
| Married | 40.6 | (±2.3) | 27.4 | (±2.3) | 6.1 | (±1.4) | 5.2 | (±1.4) |
| Not Married | 36.0 | (±2.5) | 20.5 | (±2.5) | 14.1 | (±1.9) | 4.9 | (±1.1) |
| Household Poverty Status | | | | | | | | |
| Below 100% FPL | 29.7 | (±4.4) | 15.8 | (±3.4) | 5.4 | (±2.7) | 1.7 | (±1.2) |
| At or Above 100% FPL | 37.8 | (±2.7) | 24.0 | (±2.6) | 6.4 | (±1.8) | 4.4 | (±1.6) |
| Unspecified | 42.1 | (±2.4) | 27.0 | (±2.1) | 15.2 | (±1.9) | 7.0 | (±1.2) |
| Residence | | | | | | | | |
| Urban | 40.3 | (±1.9) | 24.0 | (±1.8) | 10.8 | (±1.4) | 5.1 | (±1.0) |
| Rural | 31.7 | (±3.3) | 23.9 | (±3.3) | 6.2 | (±2.2) | 4.9 | (±1.6) |
| DHEC Region | | | | | | | | |
| Upstate | 38.8 | (±3.3) | 26.2 | (±3.3) | 11.8 | (±2.9) | 6.9 | (±2.2) |
| Pee Dee | 38.4 | (±3.2) | 25.5 | (±2.9) | 6.7 | (±1.8) | 5.6 | (±1.8) |
| Midlands | 37.2 | (±3.1) | 22.2 | (±2.7) | 7.9 | (±1.7) | 4.1 | (±1.2) |
| Low Country | 41.0 | (±3.4) | 22.2 | (±2.8) | 13.8 | (±2.6) | 3.2 | (±1.1) |

1 - To determine the ever use of tobacco, respondents were asked "Have you ever tried smoking cigars, cigarillos, or very small cigars that look like cigarettes in your entire life, even one or two puffs?", "Have you ever tried chewing tobacco, snuff, or dip, such as Skoal, Copenhagen, Grizzly, Levi Garrett, Red Man, or Day's Work, even just one time in your entire life?", "Have you ever tried smoking tobacco in a water pipe in your entire life, even one or two puffs?", "Have you ever tried snus, even just one time in your entire life?".

2 - Ninety-five percent confidence interval

3 - Data are statistically unreliable because sample size was <50. Note large confidence intervals for distribution by race/ethnicity

Table 7: Percentage of South Carolina adults who report current use of menthol cigarettes¹ and other tobacco products² (cigars, smokeless tobacco, water pipe, regular pipe, and snus), by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | Menthol Cigarettes | | Cigars | | Smokeless Tobacco | | Water Pipe | | Regular Pipe | | Snus | |
|--|--------------------|---------------------|------------|---------------|-------------------|---------------|------------|---------------|--------------|---------------|-----------------|---------------|
| | % | 95% CI ³ | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI |
| TOTAL | 53.4 | (±4.7) | 4.7 | (±0.8) | 3.6 | (±0.7) | 1.7 | (±0.5) | 1.0 | (±0.3) | 0.5 | (±0.2) |
| Gender | | | | | | | | | | | | |
| Male | 52.8 | (±6.7) | 7.4 | (±1.4) | 6.8 | (±1.4) | 2.4 | (±1.0) | 1.8 | (±0.7) | 0.9 | (±0.5) |
| Female | 54.2 | (±6.5) | 2.3 | (±0.7) | 0.7 | (±0.3) | 1.1 | (±0.5) | 0.2 | (±0.2) | 0.1 | (±0.1) |
| Age Group | | | | | | | | | | | | |
| 18-24 | 65.6 | (±14.3) | 11.0 | (±3.6) | 6.0 | (±3.1) | 5.8 | (±2.6) | 1.8 | (±1.4) | -- ⁴ | -- |
| 25-34 | 74.7 | (±9.3) | 7.1 | (±2.4) | 4.4 | (±2.2) | 3.2 | (±1.7) | 0.8 | (±0.7) | 1.0 | (±0.9) |
| 35-44 | 42.8 | (±12.2) | 5.1 | (±2.5) | 2.8 | (±1.5) | -- | -- | 1.1 | (±0.8) | -- | -- |
| 45-54 | 46.5 | (±8.4) | 3.6 | (±1.4) | 4.0 | (±1.5) | -- | -- | -- | -- | 0.7 | (±0.7) |
| 55-64 | 45.6 | (±8.2) | 2.9 | (±1.0) | 2.6 | (±1.1) | -- | -- | 0.5 | (±0.3) | -- | -- |
| 65+ | 27.8 | (±8.3) | 1.5 | (±0.5) | 2.2 | (±0.8) | 0.4 | (±0.4) | 0.6 | (±0.4) | 0.3 | (±0.3) |
| Race/Ethnicity | | | | | | | | | | | | |
| non-Hispanic White | 36.3 | (±6.1) | 4.1 | (±0.9) | 4.3 | (±1.0) | 1.3 | (±0.6) | 1.0 | (±0.4) | 0.7 | (±0.3) |
| non-Hispanic Black | 91.8 | (±3.5) | 5.1 | (±1.7) | 1.4 | (±0.6) | 2.0 | (±1.4) | 0.4 | (±0.3) | 0.2 | (±0.2) |
| non-Hispanic Asian | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| non-Hispanic American Indian & Alaska Native | 72.0 | (±26.0) | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| non-Hispanic Other | 55.7 | (±13.9) | 15.9 | (±7.4) | 4.0 | (±2.3) | 4.7 | (±4.0) | -- | -- | 0.4 | (±0.4) |
| Hispanic | 77.6 | (±21.4) | 5.0 | (±4.1) | -- | -- | 3.6 | (±3.1) | -- | -- | 0.4 | (±0.4) |
| Education | | | | | | | | | | | | |
| < High School | 59.1 | (±10.3) | 6.6 | (±3.1) | 5.3 | (±2.2) | 2.6 | (±2.5) | 1.0 | (±0.9) | 0.2 | (±0.2) |
| High School or GED | 51.6 | (±7.2) | 5.3 | (±1.5) | 4.5 | (±1.7) | 1.9 | (±0.9) | 1.2 | (±0.8) | 0.8 | (±0.5) |
| Some College | 50.0 | (±8.3) | 4.7 | (±1.3) | 2.9 | (±1.0) | 1.5 | (±0.7) | 0.9 | (±0.6) | 0.3 | (±0.3) |
| College Degree or Higher | 48.0 | (±11.0) | 2.9 | (±0.9) | 2.3 | (±0.9) | 0.9 | (±0.6) | 0.8 | (±0.5) | 0.7 | (±0.5) |
| Marital Status | | | | | | | | | | | | |
| Married | 38.9 | (±8.1) | 3.3 | (±0.9) | 3.4 | (±0.9) | 0.5 | (±0.3) | 0.7 | (±0.3) | 0.6 | (±0.3) |
| Not Married | 62.8 | (±5.4) | 6.4 | (±1.4) | 3.7 | (±1.1) | 2.9 | (±1.1) | 1.1 | (±1.1) | 0.5 | (±0.3) |

| Socio-Demographic Characteristic | Menthol Cigarettes | | Cigars | | Smokeless Tobacco | | Water Pipe | | Regular Pipe | | Snus | |
|----------------------------------|--------------------|---------------------|--------|--------|-------------------|--------|------------|--------|--------------|--------|------|--------|
| | % | 95% CI ³ | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI | % | 95% CI |
| Household Poverty Status | | | | | | | | | | | | |
| Below 100% FPL | 61.3 | (±11.7) | 4.5 | (±2.7) | 2.8 | (±1.5) | -- | -- | -- | -- | 0.2 | (±0.2) |
| At or Above 100% FPL | 45.3 | (±8.7) | 2.9 | (±0.9) | 3.3 | (±1.2) | 0.7 | (±0.4) | 0.5 | (±0.4) | 0.4 | (±0.3) |
| Unspecified | 55.9 | (±5.4) | 6.8 | (±1.3) | 4.3 | (±1.1) | 2.8 | (±0.9) | 1.6 | (±0.7) | 0.8 | (±0.4) |
| Residence | | | | | | | | | | | | |
| Urban | 52.5 | (±5.6) | 4.6 | (±0.9) | 3.2 | (±0.8) | 1.7 | (±0.5) | 0.9 | (±0.4) | 0.5 | (±0.3) |
| Rural | 55.0 | (±8.6) | 5.0 | (±2.0) | 4.7 | (±1.5) | 1.6 | (±1.6) | 0.7 | (±0.5) | 0.7 | (±0.5) |
| DHEC Region | | | | | | | | | | | | |
| Upstate | 52.7 | (±9.4) | 6.7 | (±2.1) | 5.5 | (±2.6) | 1.8 | (±1.4) | 1.4 | (±0.8) | 0.9 | (±0.6) |
| Pee Dee | 50.8 | (±8.4) | 4.6 | (±1.5) | 3.8 | (±1.4) | 1.2 | (±0.7) | 0.6 | (±0.4) | 0.5 | (±0.4) |
| Midlands | 52.3 | (±9.1) | 3.0 | (±0.9) | 2.0 | (±1.9) | 1.5 | (±0.8) | 0.4 | (±0.4) | 0.2 | (±0.2) |
| Low Country | 57.9 | (±8.6) | 4.7 | (±1.4) | 2.8 | (±1.8) | 2.2 | (±1.0) | 1.2 | (±0.8) | 0.4 | (±0.4) |

1 - Use of Menthol cigarettes was determined among current smokers and respondents were asked the question “During the past 30 days, were the cigarettes that you usually smoked menthol?” Those who responded “Yes” are coded as current users.

2 - To determine current use of other tobacco products, respondents were asked the following questions: “During the past 30 days, on how many days did you smoke cigars, cigarillos, or very small cigars that look like cigarettes?”, “During the past 30 days, on how many days did you use chewing tobacco, snuff, or dip, such as Skoal, Copenhagen, Grizzly, Levi Garrett, Red Man, or Day’s Work?”, “During the past 30 days, on how many days did you smoke tobacco in a water pipe?”, “During the past 30 days, on how many days did you smoke tobacco in a pipe other than a water pipe?”, “During the past 30 days, on how many days did you use snus?”. Those who used those tobacco products on 1 or more days are coded as current users.

3 - Ninety-five percent confidence interval.

4 - Data are statistically unreliable because sample size was <50. Note large confidence intervals for distribution by race/ethnicity.

Table 8: Percentage of South Carolina adults exposed¹ to Secondhand Smoke (SHS) at various locations – Results from the 2014-2015 SC Adult Tobacco Survey.

| Exposure to SHS | All | | Current Smokers | | Non-Smokers | |
|------------------------------------|------|-----------------------|-----------------|----------|-------------|----------|
| | % | (95% CI) ² | % | (95% CI) | % | (95% CI) |
| At Home | 10.6 | (±1.2) | 33.2 | (±4.2) | 5.4 | (±0.8) |
| In Vehicles | 16.3 | (±1.4) | 48.3 | (±4.7) | 8.7 | (±1.1) |
| In indoor or outdoor workplaces | 22.4 | (±2.1) | 38.8 | (±6.2) | 18.8 | (±2.1) |
| In indoor or outdoor public places | 27.5 | (±1.6) | 36.1 | (±4.7) | 25.4 | (±1.6) |

1 - Exposure to SHS is determined for the past 7 days.

2 - Ninety-five percent confidence interval

Table 9: Percentage of South Carolina adult current smokers¹ and adults exposed to Secondhand Smoke (SHS)² at various locations among those with chronic conditions³ – Results from the 2014-2015 SC Adult Tobacco Survey.

| Chronic condition | Current Smoker | | SHS exposure in Homes | | SHS exposure in Vehicles | | SHS exposure in Workplaces | | SHS exposure in Public Places | |
|---------------------|----------------|-----------------------|-----------------------|----------|--------------------------|----------|----------------------------|----------|-------------------------------|----------|
| | % | (95% CI) ⁴ | % | (95% CI) | % | (95% CI) | % | (95% CI) | % | (95% CI) |
| Heart Disease | 18.1 | (±4.1) | 12.0 | (±3.7) | 13.1 | (±3.8) | 20.0 | (±7.8) | 19.9 | (±4.1) |
| Cancer | 12.7 | (±4.0) | 5.0 | (±1.9) | 10.9 | (±3.4) | 13.2 | (±6.4) | 20.2 | (±5.6) |
| Diabetes | 18.1 | (±4.0) | 11.3 | (±2.6) | 13.3 | (±2.8) | 20.4 | (±8.9) | 22.9 | (±4.1) |
| Respiratory Disease | 40.5 | (±6.5) | 26.1 | (±6.1) | 26.9 | (±5.9) | 26.7 | (±11.3) | 29.5 | (±6.4) |
| Asthma | 25.9 | (±5.8) | 15.9 | (±4.2) | 24.5 | (±5.5) | 28.3 | (±6.9) | 32.0 | (±5.0) |
| High Cholesterol | 17.0 | (±2.3) | 10.5 | (±1.9) | 12.5 | (±2.0) | 20.4 | (±3.8) | 22.2 | (±2.5) |
| Hypertension | 19.1 | (±2.2) | 10.6 | (±1.7) | 13.4 | (±1.9) | 22.2 | (±3.6) | 23.0 | (±2.4) |

1 - To determine current smoking, respondents were asked two questions ‘Have you smoked at least 100 cigarettes in your entire life?’ and ‘Do you now smoke cigarettes every day, some days, or not at all? Those who smoked at least 100 cigarettes in lifetime and currently smoke every day or someday are coded as current smokers.

2 - Reported exposure to SHS is determined for the past 7 days.

3 - To determine chronic conditions/diseases, respondents were asked the following questions: “Have you ever been told by a doctor or other health professional that you have heart disease?”; “Have you ever been told by a doctor or other health professional that you have cancer, other than skin cancer?”; “Have you ever been told by a doctor or other health professional that you have diabetes, or sugar diabetes [note: gestational diabetes is excluded from this estimate]?”; “Have you ever been told by a doctor or other health professional that you have a chronic lung disease, such as emphysema, chronic bronchitis, or chronic obstructive pulmonary disease, also known as c-o-p-d?”; “Have you ever been told by a doctor or other health professional that you have asthma?”; “Have you ever been told by a doctor or other health professional that you have high cholesterol?”; “Have you ever been told by a doctor or other health professional that you have high blood pressure, or hypertension?”.

4 - Ninety-five percent confidence interval

Table 10: Percentage of South Carolina adults who reported being screened¹ for tobacco use by a health care provider during their visit in the past 12 months, overall and by chronic condition² status – Results from the 2014-2015 SC Adult Tobacco Survey.

| Characteristic | Screened for Tobacco Use | |
|--------------------------------|--------------------------|-----------------------|
| | % | (95% CI) ³ |
| TOTAL (All Respondents) | 95.0 | (±1.0) |
| Chronic Condition | | |
| Heart Disease | 96.3 | (±2.0) |
| Cancer | 95.9 | (±3.1) |
| Diabetes | 96.5 | (±1.9) |
| Respiratory Disease | 96.8 | (±2.8) |
| Asthma | 94.4 | (±3.4) |
| Cholesterol | 96.4 | (±1.3) |
| Hypertension | 95.8 | (±1.5) |

1 - To determine screening, all respondents were asked the question: “In the past 12 months, have you seen a doctor, dentist, nurse, or other health professional?” Those who responded “Yes” and were current somedays or everyday smokers were then asked the following question: “In the past 12 months, did any doctor, dentist, nurse, or other health professional advise you to quit smoking cigarettes or using any other tobacco products?” Those who responded “Yes” but were not current somedays or everyday smokers were asked the following question: “In the past 12 months, did any doctor, dentist, nurse, or other health professional ask if you smoke cigarettes or use any other tobacco products?”

2 - To determine chronic conditions/diseases, respondents were asked the following questions: “Have you ever been told by a doctor or other health professional that you have heart disease?”; “Have you ever been told by a doctor or other health professional that you have cancer, other than skin cancer?”; “Have you ever been told by a doctor or other health professional that you have diabetes, or sugar diabetes [note: gestational diabetes is excluded from this estimate]?”; “Have you ever been told by a doctor or other health professional that you have a chronic lung disease, such as emphysema, chronic bronchitis, or chronic obstructive pulmonary disease, also known as c-o-p-d?”; “Have you ever been told by a doctor or other health professional that you have asthma?”; “Have you ever been told by a doctor or other health professional that you have high cholesterol?”; “Have you ever been told by a doctor or other health professional that you have high blood pressure, or hypertension?”.

3 - Ninety-five percent confidence interval

Table 11: Percentage of South Carolina adults who reported receiving smoking cessation help from their health care provider during their visit in the past 12 months – Results from the 2014-2015 SC Adult Tobacco Survey.

| Health Care Professional advice and interventions to help quit | % | (95% CI) ¹ |
|--|------|-----------------------|
| Advised to quit smoking | 51.1 | (±4.1) |
| Advised to set a quit date | 31.0 | (±9.9) |
| Provided booklets, videos or websites | 65.3 | (±6.7) |
| Advised to use telephone Quitline or cessation counseling | 55.2 | (±7.3) |
| Prescribed NRT | 50.2 | (±7.5) |

1 - Ninety-five percent confidence interval

Table 12: Percentage of South Carolina adults who reported their quit intentions, awareness of cessation services and use of cessation services during their quit attempt – Results from the 2014-2015 SC Adult Tobacco Survey.

| Characteristic | % | (95% CI) ¹ |
|--|------|-----------------------|
| Intentions to quitting | 67.2 | (±4.2) |
| Awareness of cessation services (Quitline or other counseling services) among current smokers | 56.1 | (±4.0) |
| Use of cessation resources (Quitline, cessation program, medication or counseling) during quit attempt | 31.5 | (±5.7) |

1 - Ninety-five percent confidence interval

Table 13: Percentage of South Carolina adult current smokers who reported having quit for one or more days in the past 12 months, and sustained their abstinence – Results from the 2014-2015 SC Adult Tobacco Survey.

| Characteristic | % | (95% CI) ¹ |
|--|-------------|-----------------------|
| <i>Quit for 1 or more days in past 12 months, all current smokers</i> | 58.6 | (±5.1) |
| Everyday Smokers | 53.1 | (±6.3) |
| Someday Smokers | 76.0 | (±7.3) |
| <i>Sustained abstinence for smoking in the past 12 months</i> | 14.4 | (±4.4) |

1 - Ninety-five percent confidence interval

Table 14: Percentage of South Carolina adults who reported having seen one or more ads about not smoking, by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | TV Ads | | Radio Ads | | Print Media Ads | | Any Media Ads | |
|--|-------------|---------------------|-------------|---------------|-----------------|---------------|---------------|---------------|
| | % | 95% CI ¹ | % | 95% CI | % | 95% CI | % | 95% CI |
| TOTAL | 83.5 | (±1.4) | 47.2 | (±1.8) | 57.3 | (±1.8) | 90.6 | (±1.0) |
| Gender | | | | | | | | |
| Male | 84.4 | (±2.1) | 49.6 | (±2.7) | 57.2 | (±2.7) | 91.1 | (±1.6) |
| Female | 82.7 | (±1.9) | 45.0 | (±2.4) | 57.4 | (±2.3) | 90.2 | (±1.2) |
| Age Group | | | | | | | | |
| 18-24 | 90.0 | (±3.5) | 63.1 | (±6.4) | 60.2 | (±6.4) | 95.7 | (±2.4) |
| 25-34 | 81.4 | (±5.3) | 59.3 | (±5.5) | 58.7 | (±5.6) | 91.7 | (±3.7) |
| 35-44 | 82.9 | (±4.0) | 56.0 | (±4.9) | 56.1 | (±4.9) | 91.7 | (±2.3) |
| 45-54 | 88.0 | (±2.4) | 52.2 | (±3.9) | 56.9 | (±3.8) | 92.3 | (±2.0) |
| 55-64 | 82.1 | (±2.5) | 37.5 | (±3.2) | 58.2 | (±3.2) | 89.1 | (±2.0) |
| 65+ | 79.7 | (±2.0) | 24.6 | (±2.3) | 55.1 | (±2.6) | 86.1 | (±1.8) |
| Race/Ethnicity | | | | | | | | |
| non-Hispanic White | 81.8 | (±1.7) | 43.3 | (±2.2) | 56.4 | (±2.2) | 90.0 | (±1.1) |
| non-Hispanic Black | 90.3 | (±1.9) | 56.1 | (±3.5) | 59.5 | (±3.5) | 93.8 | (±1.6) |
| non-Hispanic Asian | 52.6 | (±25.5) | 40.3 | (±24.3) | 38.8 | (±22.8) | 70.9 | (±27.9) |
| non-Hispanic American Indian & Alaska Native | 76.7 | (±15.3) | 46.0 | (±19.8) | 57.2 | (±19.3) | 81.7 | (±14.6) |
| non-Hispanic Other | 83.4 | (±5.7) | 48.3 | (±7.7) | 54.9 | (±7.5) | 87.9 | (±5.4) |
| Hispanic | 72.5 | (±13.6) | 57.8 | (±12.8) | 64.4 | (±13.1) | 86.8 | (±11.5) |
| Education | | | | | | | | |
| < High School | 83.1 | (±5.4) | 49.6 | (±6.2) | 55.5 | (±6.1) | 92.2 | (±2.4) |
| High School or GED | 87.6 | (±1.9) | 47.8 | (±3.3) | 57.3 | (±3.2) | 92.1 | (±1.6) |
| Some College | 83.4 | (±2.5) | 48.8 | (±3.1) | 59.7 | (±3.0) | 90.6 | (±2.1) |
| College Degree or Higher | 79.3 | (±2.3) | 43.2 | (±2.8) | 56.0 | (±2.7) | 88.4 | (±1.6) |
| Marital Status | | | | | | | | |
| Married | 82.4 | (±1.9) | 43.8 | (±2.4) | 56.7 | (±2.4) | 89.5 | (±1.5) |
| Not Married | 84.8 | (±2.1) | 51.0 | (±2.7) | 58.2 | (±2.7) | 92.0 | (±1.3) |
| Household Poverty Status | | | | | | | | |
| Below 100% FPL | 80.9 | (±5.1) | 40.6 | (±5.7) | 53.8 | (±5.4) | 89.5 | (±2.4) |
| At or Above 100% FPL | 84.6 | (±2.0) | 43.6 | (±2.8) | 59.2 | (±2.8) | 89.9 | (±1.8) |
| Unspecified | 83.4 | (±1.8) | 53.6 | (±2.4) | 56.6 | (±2.4) | 91.9 | (±1.2) |
| Residence | | | | | | | | |
| Urban | 83.1 | (±1.6) | 46.8 | (±2.1) | 56.1 | (±2.1) | 90.5 | (±1.2) |
| Rural | 85.6 | (±2.7) | 48.8 | (±3.8) | 61.7 | (±3.5) | 91.4 | (±1.8) |
| DHEC Region | | | | | | | | |
| Upstate | 82.0 | (±2.8) | 48.9 | (±3.5) | 58.2 | (±3.4) | 89.3 | (±2.4) |
| Pee Dee | 88.0 | (±2.0) | 45.3 | (±3.4) | 58.3 | (±3.4) | 93.4 | (±1.5) |
| Midlands | 83.5 | (±2.7) | 47.5 | (±3.4) | 56.2 | (±3.4) | 90.7 | (±1.5) |
| Low Country | 81.8 | (±2.8) | 45.8 | (±3.6) | 56.9 | (±3.5) | 90.3 | (±2.1) |

1 - Ninety-five percent confidence interval

Table 15: Percentage of South Carolina adults who stated that the ads seen contributed to their attempt to quit, by demographics – Results from the 2014-2015 SC Adult Tobacco Survey.

| Socio-Demographic Characteristic | TV Ads | | Radio Ads | | Print Media Ads | | Any Media Ads | |
|--|-----------------|---------------------|------------|---------------|-----------------|---------------|---------------|---------------|
| | % | 95% CI ¹ | % | 95% CI | % | 95% CI | % | 95% CI |
| TOTAL | 37.6 | (±6.4) | 9.7 | (±3.9) | 12.2 | (±4.2) | 42.3 | (±6.7) |
| Gender | | | | | | | | |
| Male | 43.5 | (±9.2) | 13.7 | (±7.4) | 13.6 | (±7.4) | 49.5 | (±9.4) |
| Female | 32.2 | (±7.8) | 6.2 | (±3.2) | 11.0 | (±4.4) | 35.9 | (±8.2) |
| Age Group | | | | | | | | |
| 18-24 | 31.0 | (±14.9) | 7.9 | (±7.1) | 8.5 | (±8.5) | 39.0 | (±15.5) |
| 25-34 | 33.8 | (±14.6) | 7.7 | (±5.3) | 2.9 | (±2.9) | 36.1 | (±15.0) |
| 35-44 | 39.6 | (±15.5) | 14.0 | (±14.0) | 18.2 | (±15.7) | 42.3 | (±15.7) |
| 45-54 | 49.7 | (±11.8) | 11.2 | (±7.8) | 16.2 | (±9.4) | 54.9 | (±11.6) |
| 55-64 | 40.0 | (±11.7) | 14.0 | (±11.0) | 22.1 | (±11.5) | 46.4 | (±11.6) |
| 65+ | 33.3 | (±13.4) | 1.3 | (±1.3) | 19.8 | (±10.9) | 41.3 | (±14.5) |
| Race/Ethnicity | | | | | | | | |
| non-Hispanic White | 28.3 | (±7.6) | 4.2 | (±2.6) | 7.2 | (±3.3) | 33.9 | (±8.1) |
| non-Hispanic Black | 58.2 | (±10.8) | 21.3 | (±11.1) | 23.6 | (±11.2) | 61.7 | (±10.8) |
| non-Hispanic Asian | -- ² | -- | 64.8 | (±63.4) | 64.8 | (±63.4) | 64.8 | (±63.4) |
| non-Hispanic American Indian & Alaska Native | 33.5 | (±33.5) | 27.0 | (±27.0) | 46.8 | (±44.2) | 61.5 | (±45.8) |
| non-Hispanic Other | 53.5 | (±19.5) | 17.9 | (±16.2) | 14.8 | (±14.8) | 59.1 | (±19.2) |
| Hispanic | 20.2 | (±20.2) | -- | -- | 3.9 | (±3.9) | 21.5 | (±21.5) |
| Education | | | | | | | | |
| < High School | 45.2 | (±15.5) | 17.7 | (±12.5) | 22.7 | (±13.4) | 48.1 | (±16.3) |
| High School or GED | 40.4 | (±9.6) | 8.7 | (±4.5) | 10.7 | (±4.9) | 45.9 | (±9.8) |
| Some College | 28.1 | (±9.4) | 5.3 | (±4.0) | 6.0 | (±3.9) | 33.9 | (±10.2) |
| College Degree or Higher | 31.9 | (±13.1) | 4.1 | (±4.1) | 8.3 | (±8.2) | 36.2 | (±13.4) |
| Marital Status | | | | | | | | |
| Married | 31.9 | (±10.0) | 5.4 | (±3.7) | 7.8 | (±4.7) | 36.1 | (±10.4) |
| Not Married | 41.4 | (±8.0) | 12.3 | (±5.8) | 15.0 | (±6.0) | 46.6 | (±8.3) |
| Household Poverty Status | | | | | | | | |
| Below 100% FPL | 38.7 | (±20.1) | 13.6 | (±13.6) | 25.6 | (±17.3) | 44.3 | (±20.7) |
| At or Above 100% FPL | 41.1 | (±11.8) | 4.6 | (±3.7) | 14.4 | (±6.7) | 42.9 | (±12.0) |
| Unspecified | 34.6 | (±6.6) | 12.1 | (±4.6) | 5.6 | (±3.0) | 41.2 | (±6.9) |
| Residence | | | | | | | | |
| Urban | 34.1 | (±7.0) | 7.8 | (±3.4) | 10.4 | (±3.8) | 38.6 | (±7.3) |
| Rural | 55.5 | (±13.3) | 20.7 | (±15.6) | 20.5 | (±16.0) | 58.9 | (±12.8) |
| DHEC Region | | | | | | | | |
| Upstate | 32.7 | (±11.5) | 8.9 | (±8.9) | 17.8 | (±10.7) | 37.2 | (±12.0) |
| Pee Dee | 43.4 | (±12.5) | 13.9 | (±7.0) | 7.8 | (±5.2) | 48.4 | (±12.2) |
| Midlands | 39.7 | (±12.4) | 10.1 | (±6.3) | 9.7 | (±5.7) | 42.8 | (±13.0) |
| Low Country | 35.0 | (±11.1) | 7.1 | (±5.2) | 9.7 | (±5.4) | 42.6 | (±11.8) |

1 - Ninety-five percent confidence interval

2 - Data are statistically unreliable because sample size was <50. Note large confidence intervals for race/ethnicity.

Table 16: Percentage of South Carolina adults who mentioned that smoking is never allowed at various locations, overall and by current smoking status – Results from the 2014-2015 SC Adult Tobacco Survey.

| Smoking is never allowed in | All Respondents | | Current Smokers | | Non-Smokers | |
|-----------------------------|-----------------|-----------------------|-----------------|----------|-------------|----------|
| | % | (95% CI) ¹ | % | (95% CI) | % | (95% CI) |
| Homes | 83.7 | (±1.3) | 52.8 | (±4.7) | 91.0 | (±1.0) |
| Vehicles | 74.8 | (±1.6) | 28.7 | (±4.4) | 85.6 | (±1.4) |
| Indoor workplaces | 92.4 | (±1.5) | 85.3 | (±5.9) | 93.8 | (±1.3) |
| Outdoor workplaces | 30.5 | (±2.2) | 14.1 | (±4.1) | 34.2 | (±2.4) |

1 - Ninety-five percent confidence interval

Table 17: Percentage of South Carolina adults who thought smoking should never be allowed at various locations, overall and by current smoking status – Results from the 2014-2015 SC Adult Tobacco Survey.

| Smoking should never be allowed | All Respondents | | Current Smokers | | Non-Smokers | |
|-----------------------------------|-----------------|-----------------------|-----------------|----------|-------------|----------|
| | % | (95% CI) ¹ | % | (95% CI) | % | (95% CI) |
| At Home | 76.4 | (±1.5) | 56.9 | (±4.8) | 81.0 | (±1.4) |
| Indoor in Workplaces | 83.0 | (±1.4) | 64.5 | (±4.8) | 87.5 | (±1.2) |
| Indoors in Restaurants | 78.4 | (±1.5) | 58.6 | (±4.7) | 83.1 | (±1.4) |
| Indoors in Bars, Casinos or Clubs | 55.0 | (±1.8) | 26.2 | (±3.7) | 61.9 | (±1.8) |
| At Parks | 49.8 | (±1.8) | 32.1 | (±4.5) | 53.9 | (±1.8) |

1 - Ninety-five percent confidence interval

Table 18: Percentage of South Carolina adults who support a cigarette tax increase – Results from the 2014-2015 SC Adult Tobacco Survey.

| Characteristic | Support cigarette tax increase | |
|--------------------------------|--------------------------------|-----------------------|
| | % | (95% CI) ¹ |
| TOTAL (All Respondents) | 72.9 | (±1.6) |
| Smoking Status | | |
| Current Smokers | 52.4 | (±4.8) |
| Non Smokers | 77.9 | (±1.5) |
| DHEC Region | | |
| Upstate | 75.9 | (±2.9) |
| Midlands | 69.9 | (±3.0) |
| Pee Dee | 72.9 | (±3.0) |
| Low Country | 71.8 | (±3.3) |
| Residence | | |
| Urban | 73.0 | (±1.8) |
| Rural | 73.6 | (±3.0) |

1 - Ninety-five percent confidence interval