Hospital Infections Disclosure Act Report

Reported by: South Carolina Department of Health and Environmental Control Surgical Site Infection (SSI) Standardized Infection Ratio by Procedure

Data Collected: 01/01/2016 - 12/31/2016

Procedure	No. of Specific Procedures Performed ^a	No. of Infections	No. of Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval
Coronary Bypass Graft (Chest and Donor Incision)	364	3	2.59	1.16	0.294, 3.149
Coronary Bypass Graft (Chest Only Incision)	7	*	*	*	*
Abdominal Hysterectomy	121	0	0.63	*	*
Hip Prosthesis (Replacement)	312	4	1.80	2.22	0.705, 5.349
Knee Prosthesis (Replacement)	375	3	1.57	1.91	0.486, 5.201
Colon Surgery	242	4	5.93	0.68	0.214, 1.627

a. *= Too few procedures. Reporting on too few procedures is a risk to patient confidentiality and data stability. If less than twenty surgical procedures are performed, the SIR and number of infections will be suppressed until more procedures are performed.

Central Line Associated Blood Stream Infection (CLABSI) Standardized Infection Ratio (SIR) Data Collected: 01/01/2016 - 12/31/2016

Locationa	No. of Central Line Days ^{b,c}	No. of Infections	No. of Predicted Infections	Standardized Infection Ratio	95% Confidence Interval
All Adult Critical Care Units	7047	10	7.951	1.257	0.632, 2.219
All Adult Inpatient Wards	10013	7	9.924	0.705	0.141, 1.514
All Pediatric Critical Care Units	58	0	0.08	*	*
All Pediatric Inpatient Wards	113	1	0.11	*	*

a. The specific patient care area in which a patient is assigned while receiving care in the healthcare facility. All adult and pediatric critical care units are combined into one SIR; all adult and pediatric inpatient wards are combined into one SIR for this report.

b. Central line days are the total number of days a central line is in place for all patients in selected hospital locations.

c. * = Too few central line days. Reporting on too few central line days is a risk to confidentiality and data stability. If there are less than fifty central line days, the SIR and number of infections will be suppressed until there are more central line days to report.

Methicillin-resistant Staphylococcus aureus bloodstream infection (MRSA BSI) LabID Event Data Facility Wide Inpatient Data Collected: 01/01/2016 - 12/31/2016

Hospital Onset MRSA BSI Standardized Infection Ratio (SIR)					
No. Patient Days	nt Days No. LabID Events Predicted No. of LabID Events		SIR	95% Confidence Interval	
87081	8	7.0482401	1.135	0.527, 2.155	

a. Hospital Onset: LabID event specimen collected as an inpatient >3 days after admission to the facility (i.e., on or after day 4)

Clostridium Difficile Infections(CDI) LabID Event Data Facility Wide Inpatient Data Collected: 01/01/2016 - 12/31/2016

Hospital Onset CDI LabID Event Data					
No. Patient Days	No. of LabID Events ^a	Predicted No. of LabID Events	SIR	95% Confidence Interval	
84579	56	51.301739	1.092	0.832, 1.407	

a. Hospital Onset: LabID event specimen collected as an inpatient >3 days after admission to the facility (i.e., on or after day 4)